

UC Irvine

UC Irvine Previously Published Works

Title

Synthesis and Reduction of Bimetallic Methyl-Bridged Rare-Earth Metal Complexes, $[(C_5H_4SiMe_3)_2Ln(\mu-CH_3)]_2$ (Ln = Y, Tb, Dy).

Permalink

<https://escholarship.org/uc/item/7dj7k8f9>

Journal

ACS omega, 4(1)

ISSN

2470-1343

Authors

Dumas, Megan T
Ziller, Joseph W
Evans, William J

Publication Date

2019

DOI

10.1021/acsomega.8b02665

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed


Synthesis and Reduction of Bimetallic Methyl-Bridged Rare-Earth Metal Complexes, $[(C_5H_4SiMe_3)_2Ln(\mu-CH_3)]_2$ ($Ln = Y, Tb, Dy$)

Megan T. Dumas,¹ Joseph W. Ziller, and William J. Evans^{*,1}

Department of Chemistry, University of California, Irvine, California 92697-2025, United States


Supporting Information

ABSTRACT: The complexes $[Cp'_2Ln(\mu-CH_3)]_2$ ($Cp' = C_5H_4SiMe_3$; $Ln = Y, Tb, Dy$) were reduced to determine if these methyl-bridged complexes would form mixed valent $4f^05d^1 Ln(II)/4f^1 Ln(III)$ compounds or bimetallic $4f^05d^1 Ln(II)$ compounds containing $5d^1-5d^1$ metal–metal bonds upon reduction. Reaction of the known bridged-chloride complexes, $[Cp'_2Ln(\mu-Cl)]_2$, **1-Ln** ($Ln = Y, Tb, Dy$), with MeLi forms the bridged-methyl complexes $[Cp'_2Ln(\mu-CH_3)]_2$, **2-Ln**, which were crystallographically characterized for Tb and Dy. KC_8 reduction of **2-Ln** in the presence of 2.2.2-cryptand produced **3-Y**, **3-Tb**, and **3-Dy**, which exhibited intense dark colors and broad absorbance peaks around 400 nm with molar extinction coefficients of 1700, 2300, and 1800 $M^{-1} cm^{-1}$, respectively, which are characteristics of $Ln(II)$ ions. The dark maroon **3-Y** product had an axial electron paramagnetic resonance spectrum at 77 K ($g_1 = 1.99$, $A_1 = 17.9$ G; $g_2 = 2.00$, $A_2 = 17.7$ G) and a two-line isotropic spectrum at 273 K ($g = 1.99$, $A = 18.4$ G), which indicates that an Y(II) ion is present. Although these results are indicative of $Ln(II)$ ions present in the solution, crystallographic evidence was not obtained to establish the structure of these complexes.



INTRODUCTION

The +2 oxidation state has recently been identified in crystallographically characterizable monometallic molecular complexes of yttrium and all of the lanthanides (except radioactive Pm). This was accomplished using the coordination environment of three trimethylsilyl-substituted cyclopentadienyl ligands, as shown in eq 1.^{1–6} Previously, the +2 oxidation state in molecular species had been limited to Eu, Yb, Sm, Tm, Dy, and Nd, which had $4f^{n+1}$ electron configurations obtained by reducing a $4f^n Ln(III)$ precursor.^{7–10} Spectroscopic and structural characterizations as well as density functional theory (DFT) calculations have shown that in the trigonal coordination environment of a $[(C_5H_3RR')_3]^{-3}$ ligand set ($R = H, SiMe_3$, $R' = SiMe_3$), the new lanthanide ions exhibited a $4f^05d^1$ ground state and $4d^1$ configuration for yttrium.^{1–6} Examples of these new +2 ions are also known with other ligands including $C_5H_3(CMe_3)_2$ (Cp^{tt}),^{11,12} and $C_5H_2(CMe_3)_3$ (Cp^{ttt}),^{13,14} the tris(aryloxide) mesitylene ligand, $((Ad_MeArO)_3mes)^{3-}$,^{15,16} and $N(SiMe_3)_2$.¹⁷



$R = SiMe_3$; $M = La, Ce, Pr, Nd$
 $R = H$; $M = Y, La, Ce, Pr, Nd, Sm, Gd, Tb, Dy, Ho, Er, Tm, Lu$

The discovery of the $4f^05d^1$ electron configurations in these new ions presents the possibility of generating complexes which contain Ln–Ln bonds. Traditionally, metal–metal bonding between two lanthanides has been considered unlikely because the 4f orbitals have too limited a radial extension from the nucleus to have significant overlap. However,

electrons in d orbitals are well suited to make metal–metal bonds. Since the complexes of new $4f^05d^1 Ln(II)$ ions involve monometallic complexes with sterically crowded ligand environments, they are not ideal for placing two lanthanides in close proximity to bond.


To address this problem, reductions of $[Cp'_2Y(\mu-Cl)]_2$ and $[Cp'_2Y(\mu-H)(THF)]_2$ were explored.¹⁸ DFT calculations suggested that reduction of these bimetallic complexes could form complexes containing Y–Y bonds.¹⁸ Although spectroscopic studies, including electron paramagnetic resonance (EPR) spectroscopy and UV–vis spectroscopy, of the reductions of these complexes suggested the formation of an Y(II) ion,^{6,18,19} crystallographic evidence was elusive. In the case of complexes containing electron-deficient bridged hydride ligands, $[Cp'_2Y(\mu-H)(THF)]_2$, only trimetallic tetrahydride complexes, $[K(2.2.2-cryptand)]\{[Cp'_2Ln(\mu-H)]_3(\mu-H)\}$ ($Ln = Y, Tb, Dy$), were isolated from the reduction reactions.¹⁸ The reduction of $[Cp'_2Y(\mu-Cl)]_2$ provided an EPR spectrum at 77 K consistent with an Y(II) ion. However, this complex was extremely thermally unstable, which prohibited the collection of a room-temperature EPR spectrum, and a crystal structure could not be obtained.

To avoid the possibility of forming such trimetallic hydride-centered products, the reduction of complexes containing electron-deficient methyl-bridged ligands, $[Cp'_2Ln(\mu-CH_3)]_2$, has been explored and is reported here. Reduction of the known complex, $[Cp'_2Y(\mu-CH_3)]_2$,²⁰ is described as well as

Received: October 5, 2018

Accepted: December 17, 2018

Published: January 7, 2019

Scheme 1. Syntheses of $[\text{Cp}'_2\text{Ln}(\mu\text{-Cl})]_2$, **1-Ln**, and $[\text{Cp}'_2\text{Ln}(\mu\text{-CH}_3)]_2$, **2-Ln** $\{\text{Cp}' = [\text{C}_5\text{H}_4(\text{SiMe}_3)]\}$ 

analogous reactions of the Tb and Dy analogues, which were synthesized for this study and crystallographically characterized for definitive identification.

RESULTS AND DISCUSSION

Reaction of the known bridged-chloride complexes, $[\text{Cp}'_2\text{Ln}(\mu\text{-Cl})]_2$, **1-Ln**, with 2 equiv of methyl lithium generates the bridged-methyl complexes $[\text{Cp}'_2\text{Ln}(\mu\text{-CH}_3)]_2$, **2-Ln**, needed for this study, Scheme 1 (Ln = Y, Tb, Dy). The previously reported Y complex was identified by ^1H NMR spectroscopy,²⁰ and the paramagnetic Tb and Dy complexes were identified by X-ray crystallography, Figure 1. **2-Tb** and **2-Dy** are isomorphous with **2-Y**. Crystallographic details can be found in the Supporting Information.




Figure 1. ORTEP depiction of $[\text{Cp}'_2\text{Dy}(\mu\text{-CH}_3)]_2$, **2-Dy**, with thermal ellipsoids drawn at the 50% probability level. Hydrogen atoms have been omitted for clarity.

The crystallographic data show that the methyl-bridged complexes, **2-Ln**, have shorter Ln...Ln distances than either the chloride precursors, **1-Ln**, or the tetrahydrofuran (THF)-solvated hydrides, $[\text{Cp}'_2\text{Ln}(\mu\text{-H})(\text{THF})]_2$, Table 1.¹⁸ For example, the Tb...Tb distances in the two molecules in the unit cell of **2-Tb** were determined to be 3.5415(5) and 3.5745(4) Å compared to 4.0705(4) and 3.7041(6) Å for **1-Tb** and $[\text{Cp}'_2\text{Tb}(\mu\text{-H})(\text{THF})]_2$,¹⁸ respectively. The Ln–X–Ln angles in **2-Ln** were also determined to be the smallest when compared to the chloride and hydride analogues. These were 89.73(8) and 90.17(8)° in **2-Tb** compared to 97.29(2) and 120.7(2)° in **1-Tb** and $[\text{Cp}'_2\text{Tb}(\mu\text{-H})(\text{THF})]_2$,¹⁸ respectively. These smaller distances could


Table 1. Comparison of the Ln...Ln Distances and Ln–X–Ln Angles in **1-Ln**, **2-Ln**, and $[\text{Cp}'_2\text{Ln}(\mu\text{-H})(\text{THF})]_2$

complex	Ln...Ln distance (Å)	Ln–X–Ln angle (deg)
$[\text{Cp}'_2\text{Y}(\mu\text{-CH}_3)]_2$, 2-Y ²⁰	3.5361(9)	89.18(13)
$[\text{Cp}'_2\text{Y}(\mu\text{-Cl})]_2$, 1-Y ²¹	4.0792(9)	98.4(1)
$[\text{Cp}'_2\text{Tb}(\mu\text{-CH}_3)]_2$, 2-Tb ^a	3.5415(5)	90.17(8)
	3.5745(4)	89.73(8)
$[\text{Cp}'_2\text{Tb}(\mu\text{-H})(\text{THF})]_2$ ¹⁸	3.7041(6)	120.7(2)
$[\text{Cp}'_2\text{Tb}(\mu\text{-Cl})]_2$, 1-Tb ¹⁸	4.0705(4)	97.29(2)
$[\text{Cp}'_2\text{Dy}(\mu\text{-CH}_3)]_2$, 2-Dy ^a	3.5219(5)	90.36(8)
	3.5543(5)	89.70(8)
$[\text{Cp}'_2\text{Dy}(\mu\text{-H})(\text{THF})]_2$ ¹⁸	3.7042(3)	119.4(14)
$[\text{Cp}'_2\text{Dy}(\mu\text{-Cl})]_2$, 1-Dy ¹⁸	4.0806(5)	97.89(1)

^aUnit cell contains two molecules.

facilitate an interaction between the two rare-earth ions upon reduction by bringing the metal centers closer to each other.

Reduction of the **2-Ln** complexes with 1 equiv of KC_8 in the presence of 1 equiv of 2.2.2-cryptand (crypt) in THF produced intensely colored red-brown products, **3-Lneq 2**, typical of crystallographically characterized complexes of rare-earth ions in the +2 oxidation state, eq 1.^{2–4,6,18} The EPR spectrum of **3-Y** at 77 K has an axial signal at $g_1 = 1.99$, $A_1 = 17.9$ G; $g_2 = 2.00$, $A_2 = 17.7$ G, and the room-temperature EPR spectrum has an isotropic signal at $g = 1.99$, $A = 18.4$ G (Figure 2). These patterns are consistent with the interaction of an unpaired electron with a ^{89}Y nucleus (100% abundant $I = 1/2$) and are characteristic of Y(II) complexes (Figure 2).^{3–5,11}



The UV–vis spectra of **3-Y**, **3-Tb**, and **3-Dy**, Figure 3, show broad absorbance peaks at 420, 405, and 410 nm with molar extinction coefficients of approximately 1700, 2300, and 1800 $\text{M}^{-1} \text{cm}^{-1}$, respectively, based on the moles of metal present in the sample. The ϵ values are lower estimates that are based on the amount of **2-Ln** used to form the sample assuming complete reduction to **3-Ln**. In comparison, $[\text{K}(\text{crown})][\text{Cp}'_3\text{Y}]$ and $[\text{K}(\text{crypt})][\text{Cp}'_3\text{Y}]$ have absorptions at 530 nm ($\epsilon = 2500 \text{ M}^{-1} \text{cm}^{-1}$)⁶ and 520 nm ($\epsilon = 4500 \text{ M}^{-1} \text{cm}^{-1}$), respectively.^{3,4} The data on **3-Ln** can also be compared with those from the reductions of $[\text{Cp}'_2\text{Y}(\mu\text{-H})(\text{THF})]_2$, 794 nm ($\epsilon = 1000 \text{ M}^{-1} \text{cm}^{-1}$), $[\text{Cp}'_2\text{Tb}(\mu\text{-H})(\text{THF})]_2$, 837 nm ($\epsilon = 840 \text{ M}^{-1} \text{cm}^{-1}$), and $[\text{Cp}'_2\text{Dy}(\mu\text{-H})(\text{THF})]_2$, 715 nm ($\epsilon = 640 \text{ M}^{-1} \text{cm}^{-1}$), Figure 4.¹⁸




Figure 2. EPR spectra of 3-Y at 77 K (left; $g_1 = 1.99$, $A_1 = 17.9$ G; $g_2 = 2.00$, $A_2 = 17.7$ G) and 273 K (right; $g = 1.99$, $A = 18.4$ G).




Figure 3. UV-vis spectra of 2-Ln reduction products: 3-Y ($\lambda = 420$ nm; $\epsilon = 1700$ M⁻¹ cm⁻¹), 3-Tb ($\lambda = 405$ nm; $\epsilon = 2300$ M⁻¹ cm⁻¹), and 3-Dy ($\lambda = 410$ nm; $\epsilon = 1800$ M⁻¹ cm⁻¹).




Figure 4. UV-vis spectra of [Cp'₂Ln(μ-H)(THF)₂] reduction products: Y ($\lambda = 794$ nm; $\epsilon = 1000$ M⁻¹ cm⁻¹), Tb ($\lambda = 837$ nm; $\epsilon = 840$ M⁻¹ cm⁻¹), and Dy ($\lambda = 715$ nm; $\epsilon = 640$ M⁻¹ cm⁻¹).

The dark-colored solutions of 3-Ln maintained their color in solution at -30 °C for several days, but they did not yield crystalline products suitable for definitive characterization by X-ray diffraction. Attempts to make analogues with 18-crown-6 instead of crypt were similarly unsuccessful.

CONCLUSIONS

Although the bridged-methyl complexes [Cp'₂Ln(μ-CH₃)₂], 2-Ln, can be reduced to form dark solutions with EPR and UV-vis spectroscopic features consistent with Ln(II), isolation of crystallographically characterizable Ln(II) complexes has not been possible. These results indicate that the bis(Cp') coordination environment is not optimum for crystallizing

bimetallic complexes with a rare-earth metal in the +2 oxidation state.

EXPERIMENTAL DETAILS

All manipulations and syntheses described below were conducted with the rigorous exclusion of air and water using standard Schlenk line and glovebox techniques under an argon atmosphere. Solvents were sparged with UHP argon and dried by passage through columns containing Q-5 and molecular sieves prior to use. Elemental analyses were conducted on a Perkin-Elmer 2400 Series II CHNS elemental analyzer. UV-vis spectra were collected at 298 K using a Jasco V-670 absorption spectrometer. EPR spectra were collected using X-band frequency (9.3–9.8 GHz) on a Bruker EMX spectrometer equipped with an ER041XG microwave bridge, and the magnetic field was calibrated with DPPH ($g = 2.0036$). Infrared (IR) transmittance measurements were taken as compressed solids on a Cary 630 spectrophotometer with a diamond ATR attachment. 2.2.2-Cryptand (Sigma-Aldrich) was placed under vacuum (10^{-3} Torr) overnight before use. The solvent was removed under vacuum from a solution of methyl lithium in Et₂O (Sigma-Aldrich) to isolate it as a white powder. Anhydrous LnCl₃ (Ln = Y, Tb, Dy),²² KC₈,²³ and KC₅H₄(SiMe₃) (KCp')²⁴ were prepared according to the literature. [Cp'₂Ln(μ-Cl)]₂ (Ln = Y,²¹ Tb,¹⁸ Dy¹⁸), 1-Ln, and [Cp'₂Y(μ-CH₃)₂]₂²⁰ were prepared by a modification of the literature procedure.

[Cp'₂Dy(μ-CH₃)₂]₂, 2-Dy. A solution (-30 °C) of 1-Dy (150 mg, 0.158 mmol) in Et₂O (5 mL) was slowly added to a -30 °C slurry of MeLi (12 mg, 0.546 mmol) in Et₂O (5 mL). The total volume of the solution was increased to 20 mL with cold Et₂O. The slightly cloudy colorless solution was allowed to warm to room temperature and stirred overnight. The volatiles were removed from the solution in vacuo. The product was extracted into hexane (10 mL) and then centrifuged to remove white solids, presumably LiCl. The white solids were washed with hexane (5 mL) twice. The volatiles were removed from the hexane solution to isolate a white powder. The white powder was extracted into hexane (10 mL) again, and any insoluble materials were discarded. The volatiles were removed from the hexane solution to isolate 2-Dy as a white powder (123 mg, 0.136 mmol, 86%). X-ray quality crystals were grown from a concentrated hexane solution at -30 °C. IR: 3079(w), 2950(w), 2885(w), 1440(w), 1360(w), 1343(w), 1242(s), 1176(m), 1038(s), 905(m), 890(w), 829(s), 772(s), 751(s), 688(m). Anal. calcd for C₃₄H₅₈Dy₂Si₄: expected = C, 45.17; H, 6.47. Found: C, 41.97;

H, 6.23. The low value suggests incomplete combustion, as observed in other cases.^{1,25–28} The found CH ratio of C₃₄H_{60.1} is consistent with the formula.

[Cp⁺₂Tb(μ-CH₃)₂, 2-Tb. In a procedure analogous to the synthesis of 2-Dy, 1-Tb (150 mg, 0.161 mmol) and MeLi (11 mg, 0.482 mmol) were combined in Et₂O (20 mL) to isolate 2-Tb as a white powder (133 mg, 0.148 mmol, 92%). X-ray quality crystals were grown from a concentrated hexane solution at −30 °C. IR: 3075(w), 2951(w), 2880(w), 2814(w), 2768(w), 1711(w), 1640(w), 1604(w), 1537(w), 1442(w), 1409(w), 1359(w), 1308(w), 1242(m), 1178(m), 1130(w), 1038(s), 904(m), 888(w), 830(s), 763(s), 751(s), 687(m). Anal. calcd for C₃₄H₅₈TbSi₄: expected = C, 45.52; H, 6.52. Found: C, 41.93; H, 5.90. The low value suggests incomplete combustion, as observed in other cases.^{1,25–28} The found CH ratio of C₃₄H₅₇ is consistent with the formula.

Reduction of 2-Y. In an argon-filled glovebox, a colorless solution of 2-Y (106 mg, 0.140 mmol) and crypt (53 mg, 0.141 mmol) in THF (2 mL) was chilled to −30 °C. The solution was passed through a KC₈ column chilled to −30 °C, and a dark red-brown solution, 3-Y, resulted. EPR spectra were obtained at 77 K and room temperature. UV–vis (THF) λ_{max} = 420 nm; ε = 1700 M^{−1} cm^{−1}.

Reduction of 2-Dy. In a procedure analogous to the reduction of 2-Y, 2-Dy (198 mg, 0.219 mmol) and crypt (83 mg, 0.219 mmol) were combined in THF (2 mL) and passed through a KC₈ column to form a dark red-brown solution, 3-Dy. UV–vis (THF) λ_{max} = 414 nm; ε = 1800 M^{−1} cm^{−1}.

Reduction of 2-Tb. In a procedure analogous to the reduction of 2-Y, 2-Tb (200 mg, 0.224 mmol) and crypt (84 mg, 0.224 mmol) were combined in THF (2 mL) and passed through a KC₈ column to form a dark red-brown solution, 3-Tb. UV–vis (THF) λ_{max} = 406 nm; ε = 2300 M^{−1} cm^{−1}.

X-ray Crystallographic Data. Crystallographic information for complexes 2-Tb (1871704) and 2-Dy (1871703) are in the Supporting Information.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acsomega.8b02665.

Crystallographic details, selected bond distances and crystallographic data (CIF) for 2-Tb and 2-Dy (PDF)

Structure factors for datablocks mtd23 and mtd24, respectively (PDF) (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: wevans@uci.edu.

ORCID

Megan T. Dumas: 0000-0001-7000-2130

William J. Evans: 0000-0002-0651-418X

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the U.S. National Science Foundation for support of this research (CHE-1565776). We also thank Mikey K. Wojnar and Austin J. Ryan for help with X-ray crystallography, the UCI

Laser Facility and Professor A. S. Borovik for assistance with EPR spectroscopy.

■ REFERENCES

- (1) Hitchcock, P. B.; Lappert, M. F.; Maron, L.; Protchenko, A. V. Lanthanum Does Form Stable Molecular Compounds in the +2 Oxidation State. *Angew. Chem., Int. Ed.* **2008**, *47*, 1488–1491.
- (2) MacDonald, M. R.; Bates, J. E.; Fieser, M. E.; Ziller, J. W.; Furche, F.; Evans, W. J. Expanding Rare-Earth Oxidation State Chemistry to Molecular Complexes of Holmium(II) and Erbium(II). *J. Am. Chem. Soc.* **2012**, *134*, 8420–8423.
- (3) Macdonald, M. R.; Bates, J. E.; Ziller, J. W.; Furche, F.; Evans, W. J. Completing the Series of +2 Ions for the Lanthanide Elements: Synthesis of Molecular Complexes of Pr²⁺, Gd²⁺, Tb²⁺, and Lu²⁺. *J. Am. Chem. Soc.* **2013**, *135*, 9857–9868.
- (4) Fieser, M. E.; MacDonald, M. R.; Krull, B. T.; Bates, J. E.; Ziller, J. W.; Furche, F.; Evans, W. J. Structural, Spectroscopic, and Theoretical Comparison of Traditional vs Recently Discovered Ln²⁺ Ions in the [K(2.2.2-cryptand)][(C₅H₄SiMe₃)₃Ln] Complexes: The Variable Nature of Dy²⁺ and Nd²⁺. *J. Am. Chem. Soc.* **2015**, *137*, 369–382.
- (5) Palumbo, C. T.; Darago, L. E.; Windorff, C. J.; Ziller, J. W.; Evans, W. J. Trimethylsilyl versus Bis(trimethylsilyl) Substitution in Tris(cyclopentadienyl) Complexes of La, Ce, and Pr: Comparison of Structure, Magnetic Properties, and Reactivity. *Organometallics* **2018**, *37*, 900–905.
- (6) MacDonald, M. R.; Ziller, J. W.; Evans, W. J. Synthesis of a Crystalline Molecular Complex of Y²⁺, [(18-crown-6)K]-[(C₅H₄SiMe₃)₃Y]. *J. Am. Chem. Soc.* **2011**, *133*, 15914–15917.
- (7) Bochkarev, M. N.; Fedushkin, I. L.; Fagin, A. A.; Petrovskaya, T. V.; Ziller, J. W.; Broomhall-Dillard, R. N. R.; Evans, W. J. Synthesis and Structure of the First Molecular Thulium(II) Complex: [TmI₂(MeOCH₂CH₂OMe)₃]. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 133–135.
- (8) Bochkarev, M. N.; Fagin, A. A. A New Route to Neodymium(II) and Dysprosium (II) Iodides. *Chem. - Eur. J.* **1999**, *5*, 2990–2992.
- (9) Evans, W. J.; Allen, N. T.; Ziller, J. W. The Availability of Dysprosium Diiodide as a Powerful Reducing Agent in Organic Synthesis: Reactivity Studies and Structural Analysis of DyI₂(DME)₃ and Its Naphthalene Reduction Product. *J. Am. Chem. Soc.* **2000**, *122*, 11749–11750.
- (10) Bochkarev, M. N.; Fedushkin, I. L.; Dechert, S.; Fagin, A. A.; Schumann, H. [NdI₂(thf)₅], the First Crystallographically Authenticated Neodymium(II) Complex. *Angew. Chem., Int. Ed.* **2001**, *40*, 3176–3178.
- (11) Gun'ko, Y. K.; Hitchcock, P. B.; Lappert, M. F. Activation of a C–O bond by reaction of a tris(cyclopentadienyl)lanthanide complex with an alkali metal in dimethoxyethane (DME); crystal structures of [Nd{η-C₅H₃(SiMe₃)₂-1,3}₂(μ-OMe)₂Li(DME)] and [Ce{η-C₅H₃(SiMe₃)₂-1,3}₂(μ-OMe)₂]. *J. Organomet. Chem.* **1995**, *499*, 213–219.
- (12) Cassani, M. C.; Duncalf, D. J.; Lappert, M. F. The First Example of a Crystalline Subvalent Organolanthanum Complex: [K([18]crown-6)-(η²-C₆H₆)₂][(LaCp^{tt})₂(μ-η⁶:η⁶-C₆H₆)]·2C₆H₆(Cp^{tt} = η⁵-C₅H₃But₂-1,3). *J. Am. Chem. Soc.* **1998**, *120*, 12958–12959.
- (13) Jaroschik, F.; Nief, F.; Le Goff, X. F.; Ricard, L. Isolation of stable organodysprosium(II) complexes by chemical reduction of dysprosium(III) precursors. *Organometallics* **2007**, *26*, 1123–1125.
- (14) Jaroschik, F.; Momin, A.; Nief, F.; Le Goff, X.-F.; Deacon, G. B.; Junk, P. C. Dinitrogen Reduction and C–H Activation by the Divalent Organoneodymium Complex [(C₅H₂But₃)₂Nd(μ-I)K([18]-crown-6)]. *Angew. Chem., Int. Ed.* **2009**, *48*, 1117–1121.
- (15) La Pierre, H. S.; Scheurer, A.; Heinemann, F. W.; Hieringer, W.; Meyer, K. Synthesis and characterization of a uranium(II) monoarene complex supported by δ backbonding. *Angew. Chem., Int. Ed.* **2014**, *53*, 7158–7162.
- (16) Fieser, M. E.; Palumbo, C. T.; La Pierre, H. S.; Halter, D. P.; Voora, V. K.; Ziller, J. W.; Furche, F.; Meyer, K.; Evans, W. J.

Comparisons of lanthanide/actinide +2 ions in a tris(aryloxy)arene coordination environment. *Chem. Sci.* **2017**, *8*, 7424–7433.

(17) Ryan, A. J.; Darago, L. E.; Balasubramani, S. G.; Chen, G. P.; Ziller, J. W.; Furche, F.; Long, J. R.; Evans, W. J. Synthesis, Structure, and Magnetism of Tris(amide) $[\text{Ln}\{\text{N}(\text{SiMe}_3)_2\}_3]^{1-}$ Complexes of the Non-traditional +2 Lanthanide Ions. *Chem. - Eur. J.* **2018**, *24*, 7702–7709.

(18) Dumas, M. T.; Chen, G. P.; Hu, J. Y.; Nascimento, M. A.; Rawson, J. M.; Ziller, J. W.; Furche, F.; Evans, W. J. Synthesis and reductive chemistry of bimetallic and trimetallic rare-earth metallocene hydrides with $(\text{C}_5\text{H}_4\text{SiMe}_3)^{1-}$ ligands. *Organomet. Chem.* **2017**, *849–850*, 38–47.

(19) Corbey, J. F.; Woen, D. H.; Palumbo, C. T.; Fieser, M. E.; Ziller, J. W.; Furche, F.; Evans, W. J. Ligand Effects in the Synthesis of Ln^{2+} Complexes by Reduction of Tris(cyclopentadienyl) Precursors Including C–H Bond Activation of an Indenyl Anion. *Organometallics* **2015**, *34*, 3909–3921.

(20) Schumann, H.; Keitsch, M. R.; Demtschuk, J.; Molander, G. A. Organometallic compounds of the lanthanides: Part CIXXX. Synthesis, structure and hydrolysis of ‘moderately stable’ hydrosilylation catalysts $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ln}(\mu\text{-Me})]_2$ and $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\text{tBu})_2\text{Ln}(\mu\text{-Me})]_2$. *J. Organomet. Chem.* **1999**, *582*, 70–82.

(21) Evans, W. J.; Sollberger, M. S.; Shreeve, J. L.; Olofson, J. M.; Hain, J. H.; Ziller, J. W. Formation of bimetallic, trimetallic, and pentametallic yttrium methoxide and methoxide oxide complexes from reactions of alkali-metal methoxides with bis(cyclopentadienyl) yttrium chloride. *Inorg. Chem.* **1992**, *31*, 2492–2501.

(22) Taylor, M. D. Preparation of Anhydrous Lanthanide Halides. *Chem. Rev.* **1962**, *62*, 503–511.

(23) Bergbreiter, D. E.; Killough, J. M. Reactions of potassium-graphite. *J. Am. Chem. Soc.* **1978**, *100*, 2126–2134.

(24) Peterson, J. K.; MacDonald, M. R.; Ziller, J. W.; Evans, W. J. Synthetic Aspects of $(\text{C}_5\text{H}_4\text{SiMe}_3)_3\text{Ln}$ Rare-Earth Chemistry: Formation of $(\text{C}_5\text{H}_4\text{SiMe}_3)_3\text{Lu}$ via $[(\text{C}_5\text{H}_4\text{SiMe}_3)_2\text{Ln}]^+$ Metallocene Precursors. *Organometallics* **2013**, *32*, 2625–2631.

(25) Rabe, G. W.; Ziller, J. W. Phosphido Complexes of the Lanthanides. Synthesis and x-ray Crystal Structure Determination of a Tris(phosphido) Species of Neodymium: $\text{Nd}[\text{P}(\text{SiMe}_3)_2]_3(\text{thf})_2$. *Inorg. Chem.* **1995**, *34*, 5378–5379.

(26) Langeslay, R. R.; Fieser, M. E.; Ziller, J. W.; Furche, F.; Evans, W. J. Expanding Thorium Hydride Chemistry Through Th^{2+} , Including the Synthesis of a Mixed-Valent $\text{Th}^{4+}/\text{Th}^{3+}$ Hydride Complex. *J. Am. Chem. Soc.* **2016**, *138*, 4036–4045.

(27) Fieser, M. E.; Palumbo, C. T.; La Pierre, H. S.; Halter, D. P.; Voora, V. K.; Ziller, J. W.; Furche, F.; Meyer, K.; Evans, W. J. Comparisons of lanthanide/actinide +2 ions in a tris(aryloxy)arene coordination environment. *Chem. Sci.* **2017**, *8*, 7424–7433.

(28) Woen, D. H.; Chen, G. P.; Ziller, J. W.; Boyle, T. J.; Furche, F.; Evans, W. J. Solution Synthesis, Structure, and CO_2 Reduction Reactivity of a Scandium(II) Complex, $\{\text{Sc}[\text{N}(\text{SiMe}_3)_2]_3\}^-$. *Angew. Chem., Int. Ed.* **2017**, *56*, 2050–2053.